


IN THE CLAIMS

Please amend the claims as follows:

Claims 1-9 (Canceled).

Claim 10 (Currently Amended): A process for preparing at least one alkoxylate, which comprises:

bringing at least one alkylene oxide selected from the group consisting of ethylene oxide, propylene oxide, butylene oxide, pentylene oxide, and decene oxide into contact with at least one Guerbet alcohol ~~monofunctional linear or singly or multiply branched alcohol~~ having from 10 to 15 carbon atoms in the presence of at least one double metal cyanide compound of the general formula (I):



wherein

- M^1 is at least one metal ion selected from the group consisting of Zn^{2+} , Fe^{2+} , Fe^{3+} , Co^{3+} , Ni^{2+} , Mn^{2+} , Co^{2+} , Sn^{2+} , Pb^{2+} , Mo^{4+} , Mo^{6+} , Al^{3+} , V^{4+} , V^{5+} , Sr^{2+} , W^{4+} , W^{6+} , Cr^{2+} , Cr^{3+} , Cd^{2+} , Hg^{2+} , Pd^{2+} , Pt^{2+} , V^{2+} , Mg^{2+} , Ca^{2+} , Ba^{2+} , Cu^{2+} , La^{3+} , Ce^{3+} , Ce^{4+} , Eu^{3+} , Ti^{3+} , Ti^{4+} , Ag^+ , Rh^{2+} , Rh^{3+} , Ru^{2+} , and Ru^{3+} ;
- M^2 is at least one metal ion selected from the group consisting of Fe^{2+} , Fe^{3+} , Co^{2+} , Co^{3+} , Mn^{2+} , Mn^{3+} , V^{4+} , V^{5+} , Cr^{2+} , Cr^{3+} , Rh^{3+} , Ru^{2+} , and Ir^{3+} ;
- A and X are each, independently of one another, an anion selected from the group consisting of halide, hydroxide, sulfate, carbonate, cyanide, thiocyanate, isocyanate, cyanate, carboxylate, oxalate, nitrate, nitrosyl, hydrogensulfate, phosphate, dihydrogenphosphate, hydrogenphosphate, and hydrogencarbonate;
- L is a water-miscible ligand selected from the group consisting of alcohols, aldehydes, ketones, ethers, polyethers, esters, polyesters, polycarbonate, ureas, amides,

primary, secondary and tertiary amines, ligands having a pyridine nitrogen, nitriles, sulfides, phosphides, phosphites, phosphanes, phosphonates, and phosphates;

- k is a fraction or integer greater than or equal to 0;
- P is an organic additive;
- a, b, c, d, g and n are selected so that the compound (I) is electrically neutral,

wherein c may be equal to 0;

- e is the number of ligand molecules and is a fraction or integer greater than or equal to 0;
- f and h are each, independently of one another, a fraction or integer greater than or equal to 0; and

wherein the reaction is carried out at a temperature of from ~~130~~ 140°C to 155°C.

Claim 11 (Previously Presented): The process according to claim 10, wherein

- (1) M^1 is selected from the group consisting of Zn^{2+} , Fe^{2+} , Fe^{3+} , Co^{3+} , Ni^{2+} , Mn^{2+} , and Co^{2+} ; or
- (2) M^2 is selected from the group consisting of Fe^{2+} , Fe^{3+} , and Co^{3+} .

Claim 12 (Previously Presented): The process according to claim 10, wherein M^1 is Zn^{2+} and M^2 is Co^{3+} .

Claim 13 (Previously Presented): The process according to claim 10, wherein the at least one alkylene oxide is ethylene oxide or propylene oxide.

Claim 14 (canceled):

Claim 15 (Previously Presented): The process according to claim 10, wherein the alcohol is 2-propylheptanol or an isomer mixture thereof.

Claim 16 (Previously Presented): The process according to claim 10, wherein the sum of inert gas partial pressure and alkylene oxide partial pressure is from 1.5 bar to 6.0 bar during the induction phase.

Claim 17 (canceled)

Claim 18 (currently amended): The process according to claim 14, wherein the Guerbet alcohol is selected from the group consisting of ~~ethylhexanol~~, propylheptanol[[,]] and butyl octanol.

Claim 19 (canceled)

Claim 20 (previously presented): The process according to claim 10, wherein the reaction is carried out at a temperature of from 140°C to 150°C.

Claim 21 (canceled):